

## Strigolactone and its quintet: Signalling network of Shoot branching

Bhrundha, K<sup>1</sup>, Priyadarshini, S<sup>2</sup>, Krishna Kumar, S<sup>3</sup>

<sup>1</sup> Department of Plant Physiology, Kerala Agricultural University, Thrissur, Kerala.

<sup>2</sup> Department of Agricultural Microbiology, University of Agricultural Sciences, GKVK, Bengaluru.

<sup>3</sup> Department of Agricultural Entomology, University of Agricultural Sciences, GKVK, Bengaluru.

Corresponding Author: bhrundhakrishnan98@gmail.com

Manuscript No: KN-V4-02/017

### Introduction


As with increases in global population demand for food has increased but the arable land decreased looking for ways to raise the grain yield has become a priority. To meet this challenge new plant varieties with Ideal Plant Architecture that produce much higher yield needed to be produced. The nature and relative arrangement of each component within a plant determine its architecture which is the manifestation of a balance between the environment's extrinsic limitations and the plant's internal development processes at any given moment. Plant architectural analysis is essentially a global, multilevel and dynamic approach to plant development.

A key component of plant architecture that establishes the contact between the plant and its surroundings is shoot branching. It contributes to essential processes such as the establishment and leaf area distribution that determine light interception and photosynthesis, which in turn influence growth and yield. It is an intricate process that includes bud outgrowth and shoot extension, and these in turn respond to endogenous signals (Hormones such as Strigolactone, Auxin, Cytokinin, and Abscisic acid) and exogenous signals (Light and Nitrate).

### Hormones: Endogenous signals

In hormonal regulation, When the concentration of Auxin is higher it promotes Strigolactone (SLs), a Carotenoid-derived hormonal signal via auxin transporter PIN1 which upregulates the expression of BRC1 (BRANCHED 1) transcription factor, a key repressor for shoot branching which integrates both endogenous and exogenous inputs and mediates control of branching by regulating axillary bud potential to grow (Wang *et al.*, 2019). SL degrades the protein D53 (DWARF 53) and SMXL 6/7/8 (SUPPRESSOR OF MORE AXIL-LARY GROWTH2-LIKE) in rice and *Arabidopsis* respectively are down-regulated which in turn release the negative regulator of branching such as IPA1 (Ideal Plant Architecture1) and SPL 9/13/14/15 (SQUAMOSA PROMOTER BINDING PROTEIN-LIKE), leads to the transcriptional activation of BRC1 and inhabits the lateral growth.

Simultaneously, Auxin directly inhabits the cytokinin action that plays key roles in axillary bud initiation and outgrowth, in addition to the various process of plant growth. In the case of SLs, Cytokinin and SLs have shown antagonistic/synergistic effects on various plant's developmental processes. In rice CKX9 (CYTOKI-NIN OXIDASE/DEHYDROGENASE 9), an SL-responsive gene that encodes a cytokinin oxidase to catalyze the degradation of cytokinin and regulates the plant height and tillering (Duan *et al.*, 2019).



**Fig 1: A complex regulatory/signalling network of Shoot branching.**

### Shade light and Nitrate: exogenous signals

In Arabidopsis, FHY3 (Far-red elongated hypocotyls) and FAR1 (Far-red impaired response) are the two transcriptional factor which is essential for light signaling. Under simulated shade treatment, the accumulation of FHY3 and FAR1 decline and the transcript levels of SMXL6 and SMXL7 are down-regulated leading to increased expression of BRC1 and reduced branching (Xie *et al.*, 2020). Further, the expression of BRC 1 induces transcription of HB 21/40/53 [HOMEODOMAIN LEUCINE ZIPPER PROTEIN] homeodomain leucine zipper protein transcription factors. Together with BRC1, HB 21/40/53 enhance the gene expression NCED3 (9-CIS EPOXIC-AROTENOID DIOXIGENASE 3), which in turn leads to local accumulation of ABA and it inhibits bud/tiller outgrowth (González-Grandío *et al.*, 2017).

Nitrate (NO<sub>3</sub><sup>-</sup>) directly influences the action of SL and Cytokinin synthesis in shoot and root respectively. N limitation may affect branching by increasing SL synthesis and decreasing CK synthesis in roots (De Jong *et al.*, 2014).

### Conclusion

SL and its quintet's pathway is an important integrator of metabolic and nutritional signals in shoot branching. Tilling/Branching, a key determinant of Plant architecture enables fine-tuning of plant phenotype thus increasing yields, particularly under high-density planting conditions.

### References:

De Jong, M., George, G., Ongaro, V., Williamson, L., Willetts, B., Ljung, K. & Leyser, O. (2014). Auxin and strigolactone signaling are required for the modulation of Arabidopsis shoot branching by nitrogen supply. *Plant Physiology*, 166(1), 384-395.

Duan, J., Yu, H., Yuan, K., Liao, Z., Meng, X., Jing, Y. & Li, J. (2019). Strigolactone promotes cytokinin degradation through transcriptional activation of CYTOKININ OXIDASE/DEHYDROGENASE 9 in rice. *Proceedings of the National Academy of Sciences*, 116(28), 14319-14324.

Dun, E. A., Brewer, P. B., Gillam, E. M., & Beveridge, C. A. (2023). Strigolactones and shoot branching: what is the real hormone and how does it work?. *Plant and Cell Physiology*, 64(9), 967-983.

González-Grandío, E., Pajoro, A., Franco-Zorrilla, J.M., Tarancon, C., Immink, R.G.H. and Cubas, P. (2017) Abscisic acid signaling is controlled by a BRANCHED1/HD-ZIP I cascade in *Arabidopsis* axillary buds. *Proc. Natl. Acad. Sci. U.S.A.* 114: E245–E254.

Xie, Y., Liu, Y., Ma, M., Zhou, Q., Zhao, Y., Zhao, B. & Wang, H. (2020). *Arabidopsis* FHY3 and FAR1 integrate light and strigolactone signaling to regulate branching. *Nature Communications*, 11(1), 1955.